

Random variate generation for the first hit of a ball for the symmetric stable process in \mathbb{R}^d

Luc Devroye · John P. Nolan

Received: date / Accepted: date

Abstract We provide uniformly efficient random variate generators for a collection of distributions for the hits of the symmetric stable process in \mathbb{R}^d .

Keywords Random variate generation · Simulation · Monte Carlo method · Expected time analysis · Stable processes · Hitting times

Mathematics Subject Classification (2020) 65C10 · 65C05 · 11K45 · 68U20

1 Introduction

In this note, random variate generators that are uniformly fast in starting location are derived for a family of distributions of hits of symmetric stable processes. The motivation for this work is for use in Nolan, Audus, and Douglas (2023), where these methods are used to estimate Riesz α -capacity for general sets. More precisely, let $\{X(t); t \geq 0\}$ ($d \geq 2$) be the symmetric stable process in \mathbb{R}^d of index α with $0 < \alpha \leq 2$. When $0 < \alpha < 2$, it is a process with stationary independent increments whose continuous transition density, relative to Lebesgue measure in \mathbb{R}^d , is

$$p(t, x) = (2\pi)^{-d} \int e^{i(x, \xi) - t|\xi|^\alpha} d\xi,$$

where $x, \xi \in \mathbb{R}^d$, $t > 0$, $d\xi$ is Lebesgue measure, (x, ξ) is the inner product in \mathbb{R}^d and $|\xi|^2 = (\xi, \xi)$. We have $X(0) = x$. Define

$$T = \inf\{t \geq 0 : |X(t)| > 1\},$$

L. Devroye
School of Computer Science, McGill University, Montreal
+1(514) 849-1564
E-mail: lucdevroye@gmail.com

J. P. Nolan
Department of Mathematics and Statistics, American University, Washington DC
and Applied and Computational Mathematics Division, National Institute of Standards, Gaithersburg MD
+1(202) 885-3140
E-mail: jpnolan@american.edu

$$T^* = \inf\{t \geq 0 : |X(t)| < 1\}.$$

Thus, T and T^* are the first passage times to the exterior and interior of the unit ball, respectively. Define

$$\begin{aligned}\mu(dy) &= P\{X(T) \in dy, T < \infty\}, \quad |y| \geq 1, \\ \mu^*(dy) &= P\{X(T^*) \in dy, T^* < \infty\}, \quad |y| \leq 1.\end{aligned}$$

These describe the distributions of the hits of the unit ball when $X(0) = x$. The measures are well-known, and are both given by

$$f_x(y)dy \stackrel{\text{def}}{=} \frac{\varphi(x)}{(1 - |y|^2)^{\alpha/2} \times |x - y|^d} dy.$$

where

$$\varphi(x) = \frac{\Gamma(d/2) \sin(\pi\alpha/2) (1 - |x|^2)^{\alpha/2}}{\pi^{1+d/2}}.$$

More precisely,

$$\mu(dy) = f_x(y)dy, \quad |y| \geq 1,$$

if $0 < \alpha < 2$, $|x| < 1$, and

$$\mu^*(dy) = f_x(y)dy, |y| \leq 1,$$

if $\alpha < d$, $|x| > 1$, or if $\alpha = d = 1$, $|x| > 1$. Special cases of these results are due to Spitzer (1958) and Widom (1961). The full result, including a more detailed description of the case $d = 1 < \alpha < 2$, $|x| > 1$, is given by Blumenthal, Gettoor, and Ray (1961). For a survey and more recent results, see Kyprianou (2018).

When $\alpha = 2$, $|x| > 1$, we set $T^* = \inf\{t > 0 : |X(t)| = 1\}$, and note that $X(T^*)$ is supported on the surface of the unit ball.

In this paper, we are interested in generating a random vector Y in the unit ball $B = \{y : |y| \leq 1\}$ of \mathbb{R}^d with density proportional to $f_x(y)$ when $|x| > 1$. Figure 1 shows an example of simulated hitting points of the unit ball in \mathbb{R}^3 generated by the methods described below. Throughout the paper, $S_{d-1} = \{x \in \mathbb{R}^d : |x| = 1\}$ denotes the surface of B , and Z_d is a random variable uniformly distributed on S_{d-1} . We only deal with the case $d > 1$. We drop the dependence upon x in the notation and extend the family of distributions to include the cases $\alpha = 0$ and $\alpha = 2$. For $\alpha \in [0, 2)$, we define

$$f(y) \stackrel{\text{def}}{=} \frac{1}{(1 - |y|^2)^{\alpha/2} \times |x - y|^d},$$

which is proportional to a density on B . For $\alpha = 2$, we define the measure on the surface S_{d-1} of B that is given by the Poisson kernel; it is proportional to $|x - y|^{-d}$. This corresponds to the hit position of S_{d-1} for standard Brownian motion started at x where $|x| > 1$. While formally, f is a density for all values $\alpha \in (-\infty, 2)$, we will not be concerned here with negative values of α .

For the sake of normalization, we define $x = (\lambda, 0, 0, \dots, 0)$, where $\lambda > 1$.

Finally, we will name our algorithms for easy reference later. For the Brownian case ($\alpha = 2$), we have B0, B2, B3 and Bd, while for general $\alpha \in (0, 2)$, they are called R0, R1 and R2.




Fig. 1 A sample of $n = 5000$ hitting points of the unit ball in dimension 3 for $\alpha = 1.5$ with starting point marked in red. Points are spread throughout the ball, but more concentrated near the starting point $x = (1.5, 0, 0)$.

2 Hitting distribution for exiting the unit ball when starting at $|x| < 1$

Before focusing on simulating hitting of a ball, we discuss how the related problem of exiting a ball can be solved. When the starting point is $x = 0$, we can simulate directly the hitting distribution for the exiting the sphere problem. Recall that it also uses the density $f(y)$ and that when $x = 0$,

$$f(y) = \frac{\pi^{-(d/2+1)} \Gamma(d/2) \sin(\pi\alpha/2)}{(|y|^2 - 1)^{\alpha/2} |y|^d}, \quad |y| > 1.$$

Since this is radially symmetric, it can be simulated by $X = RZ_d$, where $R = |X|$ is the amplitude/magnitude of X and Z_d is uniform on the unit sphere S_{d-1} . Using radial symmetry, the density of R is

$$h(r) = f((r, 0, \dots, 0)) \cdot \text{Area}(S_{d-1}) \cdot r^{d-1} = \frac{2 \sin(\pi\alpha/2)}{\pi r (r^2 - 1)^{\alpha/2}}, \quad r > 1.$$

A change of variable shows that $R \stackrel{\mathcal{L}}{=} 1/\sqrt{T}$ where $T \stackrel{\mathcal{L}}{=} \text{Beta}(\alpha/2, 1 - \alpha/2)$ has density h . Surprisingly, there is no dependence on dimension d in the distribution of R .

We can also simulate the hitting distribution for the complement of the unit ball when we start at $x \neq 0$. The duality property in Riesz (1938), which is also described in Section 3 of Blumenthal, Gettoor, and Ray (1961), states that if $0 < |x| < 1$, and if $x^* = x/|x|^2$ is its spherical inverse outside the unit ball, and if $Y^* \in B$ has the hitting distribution for the ball starting from x^* , its spherical inverse $Y = Y^*/|Y^*|^2$ has the hitting distribution outside B when started at $x \in B$.

3 Warm-up: The case $\alpha = 2$ —Brownian motion

Recall that $Y = (Y_1, \dots, Y_d) = X(T^*) \in S_{d-1}$ is the point of entry of the unit ball B for Brownian motion started at $x = (\lambda, 0, 0, \dots, 0)$, $\lambda > 1$, given that Brownian motion hits B . The density of Y with respect to the uniform measure on S_{d-1} is proportional to $1/\|x-y\|^d$, where we recall that $x = (\lambda, 0, \dots, 0)$ and $y \in S_{d-1}$. As $\|x-y\| \geq \lambda - 1$, we can apply this simple rejection method:

(algorithm B0 for Brownian motion, any d)

repeat

 Generate U uniformly on $[0, 1]$, $Y = (Y_1, \dots, Y_d)$ uniform on S_{d-1}

 until $U \leq \left(\frac{(\lambda-1)^2}{\lambda^2+1-2\lambda Y_1}\right)^{d/2}$

return Y

In this algorithm, we tacitly used the fact that

$$\frac{\lambda - 1}{\|x - Y\|} = \sqrt{\frac{(\lambda - 1)^2}{\lambda^2 + 1 - 2\lambda Y_1}}.$$

The expected number of iterations grows as $((\lambda + 1)/(\lambda - 1))^d$, which makes it clear that for λ near one, a more efficient algorithm is needed. The algorithms presented below all take expected time uniformly bounded over all values of λ .

We write $W = Y_1$. A simple geometric argument shows that W has density proportional to

$$f(w) \stackrel{\text{def}}{=} \frac{(1 - w^2)^{(d-3)/2}}{(1 - w^2 + (\lambda - w)^2)^{d/2}}, \quad |w| \leq 1.$$

If Z_{d-1} denotes a uniform point on S_{d-2} , i.e., on the surface of the unit ball of \mathbb{R}^{d-1} , then we note that

$$Y \stackrel{\mathcal{L}}{=} (W, \sqrt{1 - W^2} Z_{d-1}),$$

where W and Z_{d-1} are independent. The generation of Z_{d-1} is easily achieved by taking $d - 1$ independent standard normal random variates and normalizing them to be of total Euclidean length one, see Devroye (1986), for general notions of random variate generation. We now describe how to generate W .

An inspection of the density, e.g. Figure 2, shows three regimes: for $d = 2$, it is U -shaped; for $d = 3$, it is monotonically increasing on $[-1, 1]$; and for $d > 3$, the density is unimodal, and zero at both endpoints of the interval. The cases $d = 2$ and $d = 3$ have simple explicit solutions. After presenting these, we will propose a method for $d \geq 3$ that is uniformly fast over all values of λ .

3.1 The planar case: $d = 2$.

The starting density on $[-1, 1]$ is proportional to

$$f(w) \stackrel{\text{def}}{=} \frac{1}{1 + \lambda^2 - 2\lambda w} \times \frac{1}{\sqrt{1 - w^2}}.$$




Fig. 2 The unnormalized functions f are shown for $d = 2$ (top) to $d = 6$ (bottom) for a value of $\lambda = 1.5$.

Set $\gamma = \frac{2\lambda}{1+\lambda^2}$, and note that $\gamma \in [0, 1]$. Observe that $f(w) + f(-w)$ is proportional to

$$g(w) = \frac{1}{1 - (\gamma w)^2} \times \frac{1}{\sqrt{1 - w^2}},$$

where we initially will try to generate a random variate W with density proportional to g on $[0, 1]$. Given such a W , it suffices then to replace W by $-W$ with probability $f(-W)/(f(W) + f(-W))$, i.e., with probability

$$\frac{(1 + \lambda^2)^2 - (2\lambda w)^2}{2(1 + \lambda^2)(1 + \lambda^2 + 2\lambda W)} = \frac{1 + \lambda^2 - 2\lambda W}{2(1 + \lambda^2)} = \frac{1 - \gamma W}{2}.$$

Note that $g(w) \leq h(w)$, where

$$h(w) = \frac{1}{1 - \gamma w} \times \frac{1}{\sqrt{1 - w}}.$$

The density of $Y = 1/\sqrt{1 - W}$ is proportional to

$$\frac{1}{1 + \delta y^2}, y \geq 1,$$

where $\delta = (1 - \gamma)/\gamma = (\lambda - 1)^2/2\lambda$. Thus, $R = \sqrt{\delta}Y$ has density proportional to $1/(1 + r^2)$ on $[\sqrt{\delta}, \infty)$. If U denotes a uniform $[0, 1]$ random variable, then by the

inversion method,

$$Y \stackrel{\mathcal{L}}{=} \frac{\tan\left(\arctan(\sqrt{\delta}) + U\left(\frac{\pi}{2} - \arctan(\sqrt{\delta})\right)\right)}{\sqrt{\delta}}.$$

As $W = 1 - 1/Y^2$, we can obtain a random variate from g by the rejection method by accepting W with probability

$$\frac{g(W)}{h(W)} = \frac{1 - \gamma W}{1 - (\gamma W)^2} \times \frac{\sqrt{1 - W}}{\sqrt{1 - W^2}} = \frac{1}{(1 + \gamma W)\sqrt{1 + W}}.$$

Observe that this acceptance probability is at least $1/(\sqrt{2}(1 + \gamma)) \geq 1/\sqrt{8}$. Therefore, this method is uniformly fast over all choices of $\lambda > 1$. The algorithm:

(algorithm B2 for Brownian motion, $d = 2$)

define $\gamma = \frac{2\lambda}{1 + \lambda^2}$, $\delta = (\lambda - 1)^2/2\lambda$

repeat

generate U, V , i.i.d. and uniformly on $[0, 1]$

$$Y \leftarrow \frac{\tan\left(\arctan(\sqrt{\delta}) + U\left(\frac{\pi}{2} - \arctan(\sqrt{\delta})\right)\right)}{\sqrt{\delta}}$$

Set $W = 1 - 1/Y^2$

until $V \leq \frac{1}{(1 + \gamma W)\sqrt{1 + W}}$

generate V' uniformly on $[0, 1]$

if $V' \leq \frac{1 - \gamma W}{2}$ then replace W by $-W$

return $(W, S\sqrt{1 - W^2})$, where $S = \pm 1$ is a random sign

3.2 The cubic case: $d = 3$.

Just for $d = 3$, the density of W simplifies dramatically, so that we can find a direct solution by the inversion method. We obtain that if U is uniformly distributed on $[0, 1]$ then

$$W \stackrel{\mathcal{L}}{=} \frac{\lambda}{2} + \frac{1}{2\lambda} \left(1 - \frac{1}{\left(\frac{1}{\lambda+1} + \frac{2U}{\lambda^2-1}\right)^2} \right)$$

has density proportional to

$$\frac{1}{(1 - w^2 + (\lambda - w)^2)^{3/2}}, \quad |w| \leq 1.$$

This will be called algorithm B3. Exact one-liners have been known for over two decades. See, e.g., Given, Hubbard, and Douglas (1997) and Kang, Mansfield, and Douglas (2004). These are basically equivalent to the method suggested above. As $\lambda \rightarrow \infty$, we obtain $W \stackrel{\mathcal{L}}{=} 2U - 1$, which is uniformly distributed on $[0, 1]$. This confirms Archimedes's theorem which states that a uniform point on S_2 has uniform marginals.

3.3 The general case: $d \geq 3$.

For $d > 2$, we proceed by simple rejection. Using the notation for W from above, we still use the notation f for the density of W on $[-1, 1]$ (see above). We define $g(w) = f(|w|)$, and observe that $f(w) \leq g(w)$ for all $w \in [-1, 1]$, yet $\int g \leq 2$, so rejection from g is entirely feasible. As g is symmetric about zero, it suffices to find an efficient way of generating a random variable Z with density proportional to g on $[0, 1]$, and then note that SZ has density g on $[-1, 1]$ where S is an equiprobable random sign. Define

$$\gamma = \frac{(\lambda - 1)^2}{2\lambda}.$$

We observe that $g(w)$ is proportional to

$$\frac{(1-w^2)^{(d-3)/2}(\gamma + (1-w))^{\frac{d}{2}}}{\leq} h(w) \stackrel{\text{def}}{=} \frac{(2(1-w))^{\frac{d-3}{2}}}{(\gamma + (1-w))^{\frac{d}{2}}}.$$

If H has density proportional to h on $[0, 1]$, then $T = \gamma/(1-H)$ has a density that is proportional to

$$\phi(t) = \frac{1}{\sqrt{t}(1+t)^{d/2}}, t \geq \gamma.$$

We will give a generator for T that has uniformly bounded expected time over all values of γ (and thus λ). This can be used in a simple rejection algorithm that inherits the uniform expected complexity:

(algorithm Bd for Brownian motion, $d \geq 3$)

repeat forever:

 Generate U, V uniformly on $[0, 1]$

 Generate a random sign S

 Generate T

 Set $W \leftarrow 1 - \gamma/T$

 If $U \leq \left(\frac{1+W}{2}\right)^{\frac{d-3}{2}}$

 then

 if $S = 1$

 then exit the loop

 else if $V \leq \left(\frac{\gamma+(1-W)}{\gamma+(1+W)}\right)^{\frac{d}{2}}$

 then $W \leftarrow -W$ and exit the loop

 generate Z_{d-1} uniformly on S_{d-2}

 return $(W, \sqrt{1-W^2}Z_{d-1})$

3.4 A generator for T .

There are two cases, according to whether $\gamma \geq 2/d$ or $\gamma < 2/d$. If $\gamma \geq 2/d$, we bound $\phi(t) \leq 1/(\sqrt{\gamma}(1+t)^{d/2})$. A random variate with density proportional to the dominating function is given by

$$T = (1 + \gamma)U^{-2/(d-2)} - 1,$$

where U is uniform on $[0, 1]$. Thus, one can repeat generating uniform $[0, 1]^2$ pairs (U, V) until $V \leq \sqrt{\gamma/T}$, and return T . The expected complexity is bounded from above by a function of d times $\sqrt{1+1/\gamma}$, and is therefore uniformly bounded over all $\gamma \geq 2/d$. So assume that $\gamma < 2/d$. We bound

$$\phi(t) \leq \begin{cases} \phi_1(t) = \frac{1}{\sqrt{t(1+\gamma)^{d/2}}} & \text{if } \frac{2}{d} > t \geq \gamma, \\ \phi_2(t) = \frac{1}{\sqrt{\frac{2}{d}(1+t)^{\frac{d}{2}}}} & \text{if } t \geq \frac{2}{d}. \end{cases}$$

Random variates T_1 and T_2 with densities ϕ_1 and ϕ_2 can be obtained as $\left(\sqrt{\gamma} + U \left(\sqrt{\frac{2}{d}} - \sqrt{\gamma}\right)\right)^2$ and $(1 + \frac{2}{d})U^{-2/(d-2)} - 1$, respectively, where U is uniform on $[0, 1]$. We summarize the rejection algorithm, where $p = \int_{\gamma}^{2/d} \phi_1(t) dt$ and $q = \int_{2/d}^{\infty} \phi_2(t) dt$:

(generator for T , case $\gamma < 2/d$)

$$p \leftarrow 2(\sqrt{2/d} - \sqrt{\gamma})/(1+\gamma)^{\frac{d}{2}}$$

$$q \leftarrow \sqrt{\frac{d}{2}} \frac{2}{d-2} \frac{1}{(1+\frac{2}{d})^{\frac{d-2}{2}}}$$

repeat

generate U, V, V' uniformly on $[0, 1]$

if $V' \leq \frac{p}{p+q}$

then set $T \leftarrow \left(\sqrt{\gamma} + U \left(\sqrt{\frac{2}{d}} - \sqrt{\gamma}\right)\right)^2$

Accept $\leftarrow \left[V \leq \left(\frac{1+\gamma}{1+T}\right)^{\frac{d}{2}}\right]$

else set $T \leftarrow (1 + \frac{2}{d})U^{-\frac{2}{d-2}} - 1$

Accept $\leftarrow \left[V \leq \sqrt{\frac{2}{dT}}\right]$

until Accept

return T

The probability of accepting T_1 is $E \left\{ \left(\frac{1+\gamma}{1+T_1}\right)^{d/2} \right\}$, which is greater than $1/(1 + 2/d)^{d/2}$. The latter tends to $1/e$ as $d \rightarrow \infty$. The probability of accepting T_2 is $E \left\{ \sqrt{\frac{2}{dT_2}} \right\}$, which is bounded from below by a strictly positive constant uniformly over all $d > 2$. Thus, the expected time taken by the rejection algorithm for T is uniformly bounded from above over all values of $\gamma > 0$ and $d > 2$.

4 A simple rejection algorithm when $0 < \alpha < 2$

Recalling

$$f(y) \stackrel{\text{def}}{=} \frac{1}{(1 - |y|^2)^{\alpha/2} \times |x - y|^d},$$

we see that

$$f(y) \leq \frac{1}{(1 - |y|^2)^{\alpha/2}} (\lambda - 1)^{-d}.$$

This leads to a simple rejection algorithm, as a random variable with density proportional to $(1 - |y|^2)^{-\alpha/2}$ on B can be obtained as RZ_d , where R is distributed as

$$\sqrt{\text{Beta}\left(\frac{d}{2}, 1 - \frac{\alpha}{2}\right)}.$$

Here is the rejection algorithm:

(algorithm R0)

repeat

 Generate $Q \leftarrow \text{Beta}\left(\frac{d}{2}, 1 - \frac{\alpha}{2}\right)$

 Generate U uniformly on $[0, 1]$.

 Generate Z_d uniformly on S_{d-1} .

 Set $Y \leftarrow \sqrt{Q}Z_d$.

until $U(\lambda - 1)^{-d} \leq 1/|x - Y|^d$ (where $x = (\lambda, 0, 0, \dots, 0)$)

return Y

Since $|x - Y| \leq (\lambda + 1)$, we can conservatively upper bound the expected number of iterations of this algorithm by

$$\left(\frac{\lambda + 1}{\lambda - 1}\right)^d.$$

This performance deteriorates quickly when λ approaches 1. In the next section, we construct an algorithm with uniformly bounded expected time.

5 A uniformly fast algorithm for $\alpha \in [0, 2)$

Again, we let $Y = (Y_1, \dots, Y_d) = X(T^*) \in B$ be the point of entry of the unit ball B of \mathbb{R}^d when the symmetric stable process of parameter $\alpha \in (0, 2)$ starts at $X(0) = (\lambda, 0, 0, \dots, 0)$, $\lambda > 1$, given that the process enters the ball (i.e., $T^* < \infty$). We write $W = Y_1$, and $H = \sqrt{\sum_{i=2}^d Y_i^2}$, see Figure 3. A simple geometric argument shows that (W, H) has density proportional to

$$\frac{(1 - (h^2 + w^2))^{-\frac{\alpha}{2}} h^{d-2}}{(h^2 + (\lambda - w)^2)^{d/2}}, \quad |w| \leq 1, h^2 + w^2 \leq 1, h \geq 0.$$

Given (W, H) , note that

$$Y \stackrel{\mathcal{L}}{=} (W, HZ_{d-1}),$$

where (W, H) and Z_{d-1} are independent. Therefore, we have reduced our problem to a two-dimensional one. For $d = 2$, in particular, note that Z_{d-1} is merely a random sign.




Fig. 3 Definition of the the (W, H) coordinates.

Instead of working with (W, H) , it is helpful to use coordinates (Q, R) , where

$$\begin{aligned} Q &= H^2 + W^2, \\ R &= 1 - W/\sqrt{H^2 + W^2}, \end{aligned}$$

and $(Q, R) \in [0, 1] \times [0, 2]$. Vice versa,

$$\begin{aligned} W &= (1 - R)\sqrt{Q}, \\ H &= \sqrt{2R - R^2}\sqrt{Q}. \end{aligned}$$

The joint density of (Q, R) (in terms of (q, r)) is proportional to

$$\frac{(1 - q)^{-\alpha/2} q^{(d-2)/2} (2r - r^2)^{(d-3)/2}}{(q(2r - r^2) + (\lambda - (1 - r)\sqrt{q})^2)^{d/2}}, \quad 0 \leq q \leq 1, 0 \leq r \leq 2.$$

We introduce the function $\gamma = \gamma(q, r)$ for the denominator without the exponent:

$$\gamma = q(2r - r^2) + (\lambda - (1 - r)\sqrt{q})^2.$$

Observe that $(\lambda - 1)^2 \leq \gamma \leq 1 + \lambda^2$. Thus, for $\lambda \geq 5/4$, the ratio of upper to lower bound for γ is ≤ 41 , the maximum being reached at $\lambda = 5/4$. For that case, we use rejection from a density proportional to

$$(1 - q)^{-\frac{\alpha}{2}} q^{(d-2)/2} (2r - r^2)^{(d-3)/2},$$

where the first part is a beta $(d/2, 1 - \alpha/2)$ density, and the second part is proportional to the density of two times a beta $((d-1)/2, (d-1)/2)$ random variable. Thus, the following algorithm, which can be used for all values of the parameters, uses an expected number of iterations not exceeding $41^{d/2}$ for all choices of $\alpha \in [0, 2], \lambda \geq 5/4$:

(algorithm R1)

```

repeat
  Generate  $Q \leftarrow \text{Beta} \left( \frac{d}{2}, 1 - \frac{\alpha}{2} \right)$ 
  Generate  $Q' \leftarrow \text{Beta} \left( \frac{d-1}{2}, \frac{d-1}{2} \right)$ 
  Set  $R \leftarrow 2Q'$ .
  Generate a uniform  $[0, 1]$  random variable  $U$ .
until  $U^{\frac{2}{d}} \leq \frac{(\lambda-1)^2}{\gamma(Q,R)}$ 
set  $(W, H) = ((1-R)\sqrt{Q}, \sqrt{2R-R^2}\sqrt{Q})$ 
generate a uniform point  $Z_{d-1}$  on  $S_{d-2}$ 
return  $Y \leftarrow (W, HZ_{d-1})$ 

```

This leaves us with the case $\lambda \in (1, 5/4]$. To ensure uniform speed over all these choices of λ and α , we will employ a rejection method over a partition of the space. Assume that a generic density f is bounded by a function g_k , where $\{A_k, k \geq 1\}$ is a partition of the space. Let $p_k = \int_{A_k} g_k$, $p = \sum_k p_k$. Assume furthermore that there is a constant $c > 0$ such that $\int_{A_k} f \geq c \int_{A_k} g_k$. Then the following general rejection method requires an expected number of iterations that does not exceed $1/c$:

```

repeat
  Generate integer  $K$  according to distribution  $p_k/p, k \geq 1$ .
  Generate  $X$  according to a density proportional to  $g_K$  on  $A_K$ .
  Generate  $U$  uniformly on  $[0, 1]$ .
until  $U g_K(X) \leq f(X)$ 
return  $X$ 

```

Remark 1 Straightforward evaluation of $Ug \leq f$ is numerically unstable in certain cases, so it is better to evaluate $U(g/f) \leq 1$, where g/f is algebraically simplified on each of the regions A_j .

To verify the claim, observe that $\int f = 1$, and $\sum_k \int_{A_k} g_k \leq 1/c$. We use a partition into five sets. The basic function of interest is

$$f(q, r) = \frac{\zeta(q)\rho(r)}{(\gamma(q, r))^{d/2}},$$

where

$$\begin{aligned} \zeta(q) &= (1-q)^{-\alpha/2} q^{(d-2)/2}, \\ \rho(r) &= (2r-r^2)^{(d-3)/2}, \\ \gamma(q, r) &= q(2r-r^2) + (\lambda - (1-r)\sqrt{q})^2. \end{aligned}$$




Fig. 4 Partition of the region for method $R2$ when $d = 2$. The left plot shows the partition for A_1, \dots, A_5 in the (r, q) coordinates; the right plot shows the preimage of these sets in the (x_1, x_2) coordinates.

The regions are defined as follows, se Figure 4:

$$\begin{aligned}
 A_1 &: r \geq 1/16, q \geq 1/2. \\
 A_2 &: q \leq 1/2. \\
 A_3 &: r \leq (\lambda - 1)^2, q \geq 3 - 2\lambda. \\
 A_4 &: (\lambda - 1)^2 \leq r \leq 1/16, 4r \geq (1 - q)^2. \\
 A_5 &: 1/2 \leq q \leq 3 - 2\lambda, 4r \leq (1 - q)^2.
 \end{aligned}$$

Since we employ the rejection method, it suffices to bound all three factors of $f(q, r)$ from above and below on each of the five regions. We begin with $\gamma(q, r)$:

$$\begin{aligned}
 \gamma(q, r) &= q(2r - r^2) + ((\lambda - 1) + (1 - \sqrt{q}) + r\sqrt{q})^2 \\
 &\geq q(2r - r^2) + (\lambda - 1)^2 + \left(\frac{1 - q}{2}\right)^2 + r^2q \\
 &= (\lambda - 1)^2 + \left(\frac{1 - q}{2}\right)^2 + 2rq \\
 &\geq \max\left((\lambda - 1)^2, \left(\frac{1 - q}{2}\right)^2, 2rq\right),
 \end{aligned}$$

$$\geq \begin{cases} 1/16 & \text{on } A_1 \cup A_2 \\ (\lambda - 1)^2 & \text{on } A_3 \\ r & \text{on } A_4 \\ \left(\frac{1-q}{2}\right)^2 & \text{on } A_5. \end{cases}$$

and similarly,

$$\begin{aligned} \gamma(q, r) &\leq q(2r - r^2) + ((\lambda - 1) + (1 - \sqrt{q}) + r\sqrt{q})^2 \\ &\leq 3q(2r - r^2) + 3(\lambda - 1)^2 + 3(1 - q)^2 + 3r^2q \\ &= 3(\lambda - 1)^2 + 3(1 - q)^2 + 6rq, \\ &= 3(\lambda - 1)^2 + 12\left(\frac{1-q}{2}\right)^2 + 6rq, \\ &\leq 18 \max\left((\lambda - 1)^2, \left(\frac{1-q}{2}\right)^2, 2rq\right) \end{aligned}$$

and thus,

$$\gamma(q, r) \leq \begin{cases} 12 & \text{on } A_1 \\ 8.3 & \text{on } A_2 \\ 36(\lambda - 1)^2 & \text{on } A_3 \\ 36r & \text{on } A_4 \\ 36\left(\frac{1-q}{2}\right)^2 & \text{on } A_5. \end{cases}$$

We define the upper bound used for rejection in each of the five regions as $\zeta(q)\rho(r)$ times the upper bound on $\gamma(q, r)^{-d/2}$ derived above. In a few cases, we use an even larger upper bound that increases the bound at most by a multiplicative factor that does not depend upon α or λ , and thus will not affect the claim that the method is universally fast over all $\alpha \in (0, 2)$, $\lambda \in (1, 5/4]$. The bounds are all of the form

$$f(q, r) \leq g(q, r)$$

where we observe that for $d \geq 3$,

$$\begin{aligned} f(q, r) &\leq \begin{cases} 4^d (1-q)^{-\frac{\alpha}{2}} q^{(d-2)/2} (2r - r^2)^{\frac{d-3}{2}} & \text{on } A_1 \cup A_2 \\ \frac{1}{(\lambda-1)^d} (1-q)^{-\alpha/2} q^{(d-2)/2} (2r - r^2)^{(d-3)/2} & \text{on } A_3 \\ (1-q)^{-\alpha/2} q^{(d-2)/2} r^{-3/2} (2-r)^{(d-3)/2} & \text{on } A_4 \\ 2^d (1-q)^{-d-(\alpha/2)} q^{(d-2)/2} (2r - r^2)^{(d-3)/2} & \text{on } A_5 \end{cases} \\ &\leq g(q, r) \stackrel{\text{def}}{=} \begin{cases} 4^d (1-q)^{-\alpha/2} q^{(d-2)/2} (2r - r^2)^{(d-3)/2} & \text{on } A_1 \cup A_2 \\ \frac{2^{(d-3)/2}}{(\lambda-1)^d} (1-q)^{-\alpha/2} r^{(d-3)/2} & \text{on } A_3 \\ 2^{(d-3)/2} (1-q)^{-\alpha/2} r^{-3/2} & \text{on } A_4 \\ 2^d 2^{(d-3)/2} (1-q)^{-d-(\alpha/2)} r^{(d-3)/2} & \text{on } A_5. \end{cases} \end{aligned}$$

For $d = 2$, the factor $2^{(d-3)/2}$ in the expressions dealing with A_3 , A_4 and A_5 in the definition of $g(q, r)$ should be replaced by $4/\sqrt{31}$. By inspection of each of these sets

of inequalities, it is clear that in each region, the compound upper bound on $f(q, r)$ used for rejection, divided by $f(q, r)$ is bounded by a universal constant that depends upon d but not on λ or α . Thus, the rejection method that is based on the bounds given here is uniformly fast:

Proposition 1 (a) (Speed) For fixed d , the expected number of iterations performed by algorithm R2 below is uniformly bounded over $\lambda \in (1, 5/4]$, $\alpha \in (0, 2)$. Algorithms R0 is uniformly fast over all $\lambda \geq \lambda^* > 1$ and $\alpha \in (0, 2)$, while algorithm R1 is uniformly fast over all $\lambda \geq 5/4$, $\alpha \in (0, 2)$.
 (b) (Validity) Algorithms R0 and R1 can be used for all values of the parameters. Algorithm R2 is valid for $\lambda \in (1, 5/4]$, $\alpha \in (0, 2)$.

6 Putting things together

There are two tasks left to do. First we need to compute

$$p_k = \int_{A_k} g(q, r) dq dr.$$

To facilitate computations, we call $A_0 = A_1 \cup A_2$, define $p_0 = \int_{[0,1] \times [0,2]} g(q, r)$, where g is the upper bound for A_0 extended to the entire space, and will reject all random vectors that do not fall in A_0 . This does not affect the validity of proposition 1. Define

$$p = p_0 + p_3 + p_4 + p_5.$$

The values shown below include expressions that involve the beta function $B(a, b) = \Gamma(a)\Gamma(b)/\Gamma(a+b)$, and were obtained using the identity $\int_0^2 (2r - r^2)^{(d-3)/2} dr = 2^{d-2} B((d-1)/2, (d-1)/2)$.

$$\begin{aligned} p_0 &= 4^d B\left(\frac{d}{2}, 1 - \frac{\alpha}{2}\right) \times 2^{d-2} B\left(\frac{d-1}{2}, \frac{d-1}{2}\right) \\ p_3 &= \frac{2^{(d-3)/2}}{(\lambda-1)^{\alpha/2}} \frac{2^{3-(\alpha/2)}}{(2-\alpha)(d-1)} \\ p_4 &= 2^{(d-3)/2} \frac{2^{4-\alpha/2}}{\alpha(2-\alpha)} \left((\lambda-1)^{-\alpha/2} - (1/4)^{-\alpha/2} \right) \\ p_5 &= 2^{(d-3)/2} \frac{8}{\alpha(d-1)} \left((2(\lambda-1))^{-\alpha/2} - (1/2)^{-\alpha/2} \right). \end{aligned}$$

For $d = 2$, the factor $2^{(d-3)/2}$ in the expressions for p_3, p_4 and p_5 should be replaced by $4/\sqrt{31}$.

On each A_k , we need to show how to generate a random pair (Q, R) with density proportional to g . Except for A_4 and A_5 , this is quite straightforward, as we will see below.

The full algorithm:

(algorithm R2; $\lambda \in (1, 5/4]$)

repeat

 Generate a random integer K with $P\{K = k\} = p_k/p, k \in \{0, 3, 4, 5\}$.

 Generate a random pair $(Q, R) \in A_K$ with density proportional to g on A_K .

 Generate U uniformly on $[0, 1]$.

until $Ug(Q, R) \leq f(Q, R)$ and either $K > 0$ or $(K = 0, Q \leq 1/2)$ or
 $(K = 0, R \geq 1/16)$

set $W \leftarrow (1 - R)\sqrt{Q}$

set $H \leftarrow \sqrt{2R - R^2}\sqrt{Q}$

generate Z_{d-1} uniformly on S_{d-2}

return $Y \leftarrow (W, HZ_{d-1})$

The individual generators for g are as follows, where V_1 and V_2 denote independent uniform $[0, 1]$ random variables:

- (for A_0)** Generate $Q \leftarrow \text{Beta}(\frac{d}{2}, 1 - \frac{\alpha}{2})$
 Generate $R \leftarrow 2 \text{Beta}(\frac{d-1}{2}, \frac{d-1}{2})$
- (for A_3)** Generate U, V uniformly on $[0, 1]$
 Compute $Q \leftarrow 1 - 2(\lambda - 1)V^{\frac{2}{2-\alpha}}$
 Compute $R \leftarrow (\lambda - 1)^2 U^{\frac{2}{d-1}}$
- (for A_4)** Generate U, V uniformly on $[0, 1]$
 Set $\Delta \leftarrow ((\lambda - 1)^{-\alpha/2} - (1/4)^{-\alpha/2})$
 Compute $R \leftarrow ((1/4)^{-\alpha/2} + U\Delta)^{-\frac{4}{\alpha}}$
 Compute $Q \leftarrow 1 - \sqrt{4RV}^{\frac{2}{2-\alpha}}$
- (for A_5)** Generate U, V uniformly on $[0, 1]$
 Set $\Delta \leftarrow ((2(\lambda - 1))^{-\alpha/2} - (1/2)^{-\alpha/2})$
 Compute $Q \leftarrow 1 - ((1/2)^{-\alpha/2} + V\Delta)^{-\frac{2}{\alpha}}$
 Compute $R \leftarrow \frac{(1-Q)^2}{4} U^{\frac{2}{d-1}}$

7 Practical considerations

These algorithms have been coded using the open source R language, see R Core Team (2023). The Figures 5 and 6 show the hitting locations of the unit ball in the plane for varying values of α and λ .

We compared the simple rejection algorithm R0 with the uniformly fast algorithms R1 and R2. The timing shown in Table 1 shows that the performance of R0 deteriorates quickly as λ gets close to one. Furthermore, method R1 worsens with the dimension. We should point out that neither method is uniformly bounded in the dimension d . For one thing, any algorithm should take time at least linearly increasing with d .

The methods described above assume a starting point on the first axis. For a general starting point x , first rotate this point to the x_1 axis, e.g. $x \rightarrow x^* \stackrel{\text{def}}{=} (|x|, 0, \dots, 0)$. Then apply the algorithms given above with starting point x^* to produce an output y^* ,




Fig. 5 First hitting locations on the unit ball starting from $(1, 2, 0)$ for varying α in dimension $d = 2$. When $\alpha = 2$, the locations are on the surface. When $\alpha < 2$, the points are on the interior and get more uniform as α decreases to 0.




Fig. 6 First hitting locations on the unit ball starting from $(\lambda, 0)$, where λ varies for fixed $\alpha = 1.5$ in dimension $d = 2$.

n	d	λ	simple rejection	uniform bound
100000	2	1.5	0.0648	0.6170
100000	2	1.25	0.1206	0.1187
100000	2	1.1	0.3620	0.1850
100000	2	1.01	7.8300	0.0912
1000	2	1.001	210.6600	0.0714
100000	3	1.5	0.1721	0.1618
100000	3	1.25	0.5636	0.5138
100000	3	1.1	3.5275	0.5900
1000	3	1.01	675.72	0.1800
10	3	1.001	261175	0.1157
100000	4	1.5	0.4796	0.4277
100000	4	1.25	2.6138	2.2930
100000	4	1.1	36.0969	2.1580
100	4	1.01	65572.40	0.4516
1	4	1.001	∞	0.2050
100000	5	1.5	1.4049	1.2400
100000	5	1.25	12.5658	11.118
100000	5	1.1	374.1918	8.6331
10	5	1.01	4067144	1.4519
1	5	1.001	∞	0.5021

Table 1 Timing in milliseconds per random vector for the two methods with $\alpha = 1.1$. Simple rejection refers to algorithm R0 in the text. Uniform bound refers to the algorithms R2 (for $1 < \lambda \leq 1.25$) and R1 (for $\lambda \geq 1.25$). The sample size n was 100,000 for all entries under “uniform bound”; the figures given above for n are for R0 only. The time value of ∞ refers to a simulation that did not halt within eight hours for a single variate.

and then reverse the above rotation to get the final y . This rotation back to the original direction is accomplished by using d Given’s rotations.

8 The work ahead

While the algorithm above is uniformly fast over all $\lambda > 1$, $\alpha \in [0, 2)$, it is not uniformly fast over all dimensions d . Thus an improvement in that respect is desirable.

It would be quite interesting to develop an algorithm that can efficiently generate the pair (X, T) , where X is the location of entry in the unit ball and T is the time of entry. For the Brownian case ($\alpha = 2$), the joint distribution is, e.g., given in Uchiyama (2016).

References

- Blumenthal RM, Gettoor RK, Ray DB (1961) On the distribution of first hits for the symmetric stable processes. *Transactions of the American Mathematical Society* 99(3):540–554
- Devroye L (1986) *Non-Uniform Random Variate Generation*. Springer Verlag, N. Y.
- Given JA, Hubbard JB, Douglas JF (1997) A first-passage algorithm for the hydrodynamic friction and diffusion-limited reaction rate of macromolecules. *Journal of Chemical Physics* 106(9):3761–3771
- Kang EH, Mansfield ML, Douglas JF (2004) Numerical path integration technique for the calculation of transport properties of proteins. *Phys Rev E* 69:031918
- Kyprianou AE (2018) Stable processes, self-similarity and the unit ball. *ALEA, Lat Am J Probab Math Stat* 15:617–690
- Nolan JP, Audus D, Douglas J (2023) Computation of α -capacity C_α of general sets in \mathbb{R}^d using stable random walks. Preprint
- R Core Team (2023) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria
- Riesz M (1938) Intégrales de Riemann-Liouville et potentiels. *Acta Sci Math Szeged* 9:1–42
- Spitzer F (1958) Some theorems concerning 2-dimensional brownian motion. *Transactions of the AMS* 87:187–197
- Uchiyama K (2016) Density of space-time distribution of brownian first hitting of a disc and a ball. *Potential Analysis* 44:497–541
- Widom H (1961) Stable processes and integral equations. *Transactions of the American Mathematical Society* 98(3):430–449